



ENSO Cycle: Recent Evolution, Current Status and Predictions

**Update prepared by
Climate Prediction Center / NCEP
18 August 2008**



Outline

- Overview
- Recent Evolution and Current Conditions
- Oceanic Niño Index (ONI) – **“Revised November 2007”**
- Pacific SST Outlook
- U.S. Seasonal Precipitation and Temperature Outlooks
- Summary




Summary

- **ENSO-neutral conditions are present in the equatorial Pacific Ocean.**
- **Equatorial SSTs in the central Pacific Ocean have returned to near-average, while positive SST anomalies continue in the eastern Pacific.**
- **Aspects of the atmospheric circulation and pattern of tropical convection reflect a lingering La Niña signal, particularly over the western and central Pacific.**
- **Based on recent SST trends and model forecasts, ENSO-neutral conditions are expected to continue through the Northern Hemisphere Fall 2008.**



Recent Evolution of Equatorial Pacific SST Departures (°C)



Since February 2008, negative sea surface temperature anomalies have weakened over the equatorial Pacific Ocean, and positive anomalies expanded westward into the east-central equatorial Pacific Ocean.





Niño Region SST Departures (°C)

Recent Evolution

The latest weekly SST departures are:


- Niño 4 -0.2°C
- Niño 3.4 0.3°C
- Niño 3 0.8°C
- Niño 1+2 1.6°C





SST Departures (°C) in the Tropical Pacific During the Last 4 Weeks


During the last month, equatorial Pacific SSTs were below-average near the Date Line, and more than 0.5°C above-average east of 140°W.





Global SST Departures (°C)


Average SST Anomalies
20 JUL 2008 – 16 AUG 2008




Equatorial SSTs remained below-average near the Date Line and in the eastern Indian Ocean, and above-average in the eastern Pacific and across the Atlantic Ocean. Positive anomalies covered much of the North Atlantic and western North Pacific Oceans.



Weekly SST Departures (°C) for the Last Four Weeks




- During late July through mid August 2008, small negative SST departures remained slightly below average near the date line and above average in the eastern Pacific.
- Over the last month, the change in SST departures was slightly positive across much of the equatorial Pacific Ocean.





Upper-Ocean Conditions in the Eq. Pacific

Cold Episodes ↓
Warm Episodes ↑




- The basin-wide equatorial upper ocean (0-300 m) heat content is **greatest** prior to and during the early stages of a Pacific **warm** (El Niño) episode (compare top 2 panels) and **least** prior to and during the early stages of a **cold** (La Niña) episode.
- The slope of the oceanic thermocline is least (greatest) during warm (cold) episodes.
- Current values of the upper-ocean heat anomalies (slightly positive) and the thermocline slope index (near zero) indicate ENSO-neutral.

The monthly thermocline slope index represents the difference in anomalous depth of the 20°C isotherm between the western Pacific (160°E-150°W) and the eastern Pacific (90°-140°W).



Central & Eastern Pacific Upper-Ocean (0-300 m) Weekly Heat Content Anomalies


EQ. Upper-Ocean Heat Anoms. (deg C) for 180-100W



The upper ocean heat content was below-average across the eastern half of the equatorial Pacific Ocean between January 2007 and March 2008, above-average from early April 2008 through mid-July 2008, and returned to near-average during the past few weeks.




Sub-Surface Temperature Departures (°C) in the Equatorial Pacific



Longitude

- During mid June – early August 2008, the positive sub-surface temperature anomalies weakened at thermocline depth (dashed black line) across the equatorial Pacific.
- The most recent period (below) shows negative temperature anomalies at thermocline depth have strengthened in the central Pacific.




Most recent pentad analysis




Tropical OLR and Wind Anomalies During the Last 30 Days

OLR Anomalies
17 JUL 2008 to 11 AUG 2008




Positive OLR anomalies (suppressed convection and precipitation, red shading) persisted over the central and western equatorial Pacific and over the eastern Indian Ocean. Weak negative OLR anomalies were observed over parts of Indonesia.

CDAS 850-hPa Wind Anoms
17 JUL 2008-15 AUG 2008



Low-level (850-hPa) easterly wind anomalies persisted over the western equatorial Pacific Ocean. Low-level westerly anomalies continued in the eastern Pacific Ocean.

CDAS 200-hPa Wind Anoms
17 JUL 2008-15 AUG 2008




At 200-hPa, westerly wind anomalies continued in the western and central Pacific Ocean.


Atmospheric Circulation over the North Pacific & North America During the Last 60 Days

200-hPa Wind


17 JUN 2008-01 JUL 2008




02 JUL 2008-16 JUL 2008



17 JUL 2008-31 JUL 2008




01 AUG 2008-15 AUG 2008




500-hPa Height & Anoms.


17 JUN 2008-01 JUL 2008




02 JUL 2008-16 JUL 2008



17 JUL 2008-31 JUL 2008




01 AUG 2008-15 AUG 2008




925-hPa Temp. Anoms. (°C)


17 JUN 2008-01 JUL 2008




02 JUL 2008-16 JUL 2008



17 JUL 2008-31 JUL 2008



01 AUG 2008-15 AUG 2008




During mid June – mid August, the atmospheric circulation featured weak, anomalous ridging over western North America and a downstream trough over the eastern United States. During this period, the anomalous ridge contributed to the persistence of above-average temperatures over the western US and the anomalous trough led to average- to below- average temperatures in the central or eastern US.




U.S. Temperature and Precipitation Departures During the Last 30 and 90 Days

Last 30 Days

30-day (ending 17 Aug 2008) % of average precipitation




30-day (ending 16 Aug 2008) temperature departures (degree C)




Last 90 Days

90-day (ending 17 Aug 2008) % of average precipitation



90-day (ending 16 Aug 2008) temperature departures (degree C)






Intraseasonal Variability

- **Intraseasonal variability in the atmosphere (wind and pressure), which is often related to the Madden-Julian Oscillation (MJO), can significantly impact surface and subsurface conditions across the Pacific Ocean.**
- **Related to this activity**
 - **significant weakening of the low-level easterly winds usually initiates an eastward-propagating oceanic Kelvin wave.**
 - **Several Kelvin waves have occurred during the last year (see next slide).**



Weekly Heat Content Evolution in the Equatorial Pacific




- During July- February 2007, negative subsurface temperature anomalies persisted across the central and eastern equatorial Pacific. The upper-ocean heat content was affected by weak oceanic Kelvin wave activity during the period.

- Negative heat content anomalies have developed in the central equatorial Pacific, while positive anomalies have weakened in the eastern equatorial Pacific.

- Oceanic Kelvin waves have alternating warm and cold phases. The warm phase is indicated by dashed lines. Down-welling and warming occur in the leading portion of a Kelvin wave, and up-welling and cooling occur in the trailing portion.



Low-level (850-hPa) Zonal (east-west) Wind Anomalies (m s^{-1})



Westerly wind anomalies
(orange/red shading).

Easterly wind anomalies (blue
shading).


From May- June 2008, intraseasonal (MJO) activity weakened the easterly anomalies across the central equatorial Pacific (dashed oval in figure).

Low-level (850-hPa) easterly wind anomalies have persisted since January 2007 over the equatorial Pacific between 150°E and 150°W.

Low-level westerly wind anomalies have dominated in the eastern Pacific (east of 150°W) since February 2008.



200-hPa Velocity Potential Anomalies (5°N-5°S)



Positive anomalies (brown shading) indicate unfavorable conditions for precipitation.


Negative anomalies (green shading) indicate favorable conditions for precipitation.

Moderate-to-strong MJO activity was present during October 2007 – February 2008 and during mid-May – mid-June 2008.

Recently, MJO activity has been weak.



Outgoing Longwave Radiation (OLR) Anomalies



Drier-than-average conditions (orange/red shading)

Wetter-than-average conditions (blue shading)

Since February 2007, convection has been suppressed across the central equatorial Pacific Ocean.

Convection has occasionally been enhanced over the western equatorial Pacific and central Indian Ocean.



Oceanic Niño Index (ONI)

- The ONI is based on SST departures from average in the Niño 3.4 region, and is a principal measure for monitoring, assessing, and predicting ENSO.
- Defined as the three-month running-mean SST departures in the Niño 3.4 region. Departures are based on a set of improved homogeneous historical SST analyses (Extended Reconstructed SST – **ERSST.v3**). The SST reconstruction methodology is described in Smith et al., 2007, *J. Climate*, *in press*.
- Used to place current events into a historical perspective
- NOAA's operational definitions of El Niño and La Niña are keyed to the ONI index.



NOAA Operational Definitions for El Niño and La Niña

El Niño: characterized by a *positive* ONI greater than or equal to $+0.5^{\circ}\text{C}$.

La Niña: characterized by a *negative* ONI less than or equal to -0.5°C .


By historical standards, to be classified as a full-fledged El Niño or La Niña episode, these thresholds must be exceeded for a period of at least 5 consecutive overlapping 3-month seasons.

CPC considers El Niño or La Niña conditions to occur when the monthly Niño3.4 SST departures meet or exceed $\pm 0.5^{\circ}\text{C}$ along with consistent atmospheric features.



ONI (°C): Evolution since 1950

The most recent ONI value (May – July 2008) is **-0.4°C**.



El Niño
neutral
La Niña



Historical El Niño and La Niña Episodes

Based on the ONI computed using ERSST.v3

Highest		Lowest	
<u>El Niño</u>	<u>ONI Value</u>	<u>La Nina</u>	<u>ONI Value</u>
JAS 1951 - NDJ 1951/52	0.7	ASO 1949 – FMA 1951	-1.8
MAM 1957 – MJJ 1958	1.6	MAM 1954 – DJF 1956/57	-2.0
JJA 1963 – DJF 1963/64	1.0	MAM 1964 – JFM 1965	-1.1
MJJ 1965 – MAM 1966	1.6	NDJ 1967/68 – MAM 1968	-0.9
OND 1968 – MJJ 1969	1.0	JJA 1970 – DJF 1971/72	-1.4
ASO 1969 – DJF 1969/70	0.7	AMJ 1973 – JJA 1974	-2.0
AMJ 1972 – FMA 1973	2.1	ASO 1974 – AMJ 1976	-1.8
ASO 1976 – JFM 1977	0.8	SON 1984 – ASO 1985	-1.1
ASO 1977 - JFM 1978	0.8	AMJ 1988 – AMJ 1989	-2.0
AMJ 1982 – MJJ 1983	2.3	ASO 1995 – FMA 1996	-0.8
ASO 1986 – JFM 1988	1.7	JJA 1998 – MJJ 2000	-1.7
AMJ 1991 – JJA 1992	1.8	SON 2000 – JFM 2001	-0.7
JJA 1994 – FMA 1995	1.3	JAS 2007 – AMJ 2008	-1.5
AMJ 1997 – MAM 1998	2.5		
AMJ 2002 – FMA 2003	1.5		
JJA 2004 – JFM 2005	0.9		
JAS 2006 - DJF 2006/07	1.2		

NOTE:

After upgrading the ocean analysis to ERSST.v3, the following weak ENSO episodes no longer meet the NOAA criteria for an ENSO episode:

El Niño:
FMA 1993-JJA 1993

La Nina:
ASO 1961-MAM 1962
ASO 1983- DJF 1983/84



Historical Pacific warm (red) and cold (blue) episodes based on a threshold of +/- 0.5 °C for the Oceanic Nino Index (ONI) [3 month running mean of ERSST.v3 SST anomalies in the Nino 3.4 region (5N-5S, 120-170W)], calculated with respect to the 1971-2000 base period. For historical purposes El Niño and La Niña episodes are defined when the threshold is met for a minimum of 5 consecutive over-lapping seasons.

Year	DJF	JFM	FMA	MAM	AMJ	MJJ	JJA	JAS	ASO	SON	OND	NDJ
1950	-1.7	-1.5	-1.4	-1.4	-1.3	-1.2	-0.9	-0.8	-0.8	-0.8	-0.9	-1.0
1951	-1.1	-0.9	-0.7	-0.4	-0.2	0.1	0.3	0.5	0.6	0.7	0.7	0.6
1952	0.3	0.2	0.1	0.1	0.0	-0.2	-0.3	-0.3	-0.1	-0.2	-0.2	-0.1
1953	0.1	0.3	0.4	0.4	0.5	0.4	0.4	0.4	0.4	0.3	0.3	0.2
1954	0.3	0.2	-0.2	-0.6	-0.8	-0.8	-0.8	-1.1	-1.2	-1.1	-1.1	-1.0
1955	-1.0	-0.9	-0.9	-1.0	-1.1	-1.0	-1.0	-1.0	-1.4	-1.8	-2.0	-1.7
1956	-1.2	-0.7	-0.6	-0.6	-0.5	-0.5	-0.6	-0.8	-0.8	-0.9	-0.8	-0.7
1957	-0.5	-0.1	0.3	0.6	0.7	0.9	0.9	0.9	0.9	0.9	1.2	1.5
1958	1.7	1.5	1.1	0.7	0.5	0.5	0.4	0.2	0.0	0.0	0.2	0.4
1959	0.4	0.5	0.4	0.2	0.1	-0.2	-0.4	-0.5	-0.4	-0.3	-0.2	-0.3
1960	-0.3	-0.3	-0.3	-0.1	-0.1	-0.1	0.0	0.0	0.0	-0.2	-0.2	-0.2
1961	-0.1	-0.2	-0.2	-0.1	0.1	0.2	0.1	-0.3	-0.6	-0.6	-0.5	-0.4
1962	-0.5	-0.5	-0.4	-0.5	-0.4	-0.3	-0.2	-0.3	-0.4	-0.6	-0.7	-0.7
1963	-0.6	-0.3	0.0	0.1	0.1	0.3	0.7	0.9	0.9	0.9	1.0	1.0
1964	0.9	0.4	0.0	-0.5	-0.7	-0.7	-0.7	-0.8	-1.0	-1.1	-1.1	-1.0
1965	-0.8	-0.5	-0.2	0.0	0.3	0.7	1.0	1.3	1.5	1.6	1.6	1.5
1966	1.2	1.1	0.8	0.5	0.3	0.2	0.2	0.0	-0.2	-0.2	-0.3	-0.3
1967	-0.4	-0.5	-0.6	-0.5	-0.2	0.0	0.0	-0.2	-0.4	-0.5	-0.4	-0.5
1968	-0.7	-0.8	-0.8	-0.7	-0.4	0.0	0.3	0.3	0.3	0.4	0.7	0.9
1969	1.0	1.0	0.9	0.8	0.6	0.5	0.4	0.4	0.6	0.7	0.7	0.6
1970	0.5	0.3	0.2	0.1	0.0	-0.3	-0.6	-0.7	-0.7	-0.7	-0.8	-1.1
1971	-1.3	-1.4	-1.2	-0.9	-0.8	-0.8	-0.8	-0.8	-0.8	-0.9	-1.0	-0.9
1972	-0.7	-0.3	0.0	0.3	0.6	0.8	1.1	1.4	1.6	1.8	2.1	2.1
1973	1.8	1.2	0.5	0.0	-0.5	-0.8	-1.0	-1.2	-1.4	-1.7	-1.9	-2.0
1974	-1.8	-1.6	-1.2	-1.1	-0.9	-0.7	-0.5	-0.4	-0.5	-0.7	-0.8	-0.7
1975	-0.6	-0.6	-0.7	-0.8	-0.9	-1.1	-1.3	-1.3	-1.5	-1.6	-1.7	-1.7



Historical Pacific warm (red) and cold (blue) episodes based on a threshold of +/- 0.5 °C for the Oceanic Nino Index (ONI) [3 month running mean of ERSST.v3 SST anomalies in the Nino 3.4 region (5N-5S, 120-170W)], calculated with respect to the 1971-2000 base period. For historical purposes El Niño and La Niña episodes are defined when the threshold is met for a minimum of 5 consecutive over-lapping seasons.

Year	DJF	JFM	FMA	MAM	AMJ	MJJ	JJA	JAS	ASO	SON	OND	NDJ
1976	-1.6	-1.2	-0.9	-0.6	-0.5	-0.2	0.1	0.3	0.6	0.8	0.8	0.8
1977	0.6	0.5	0.3	0.2	0.2	0.4	0.4	0.4	0.5	0.7	0.8	0.8
1978	0.8	0.5	0.0	-0.3	-0.4	-0.3	-0.3	-0.4	-0.4	-0.3	-0.2	-0.1
1979	-0.1	0.0	0.1	0.2	0.1	0.0	0.1	0.2	0.3	0.5	0.5	0.6
1980	0.5	0.4	0.3	0.2	0.3	0.3	0.2	0.0	-0.1	0.0	0.0	0.0
1981	-0.2	-0.4	-0.4	-0.3	-0.2	-0.3	-0.3	-0.3	-0.2	-0.1	-0.1	0.0
1982	0.0	0.1	0.2	0.4	0.7	0.7	0.8	1.0	1.5	1.9	2.2	2.3
1983	2.3	2.1	1.6	1.3	1.0	0.7	0.3	-0.1	-0.5	-0.7	-0.9	-0.7
1984	-0.4	-0.2	-0.2	-0.3	-0.4	-0.4	-0.3	-0.2	-0.2	-0.6	-0.9	-1.1
1985	-1.0	-0.9	-0.8	-0.8	-0.8	-0.6	-0.6	-0.5	-0.6	-0.4	-0.4	-0.4
1986	-0.5	-0.5	-0.3	-0.2	-0.1	0.0	0.2	0.4	0.6	0.9	1.0	1.2
1987	1.2	1.3	1.2	1.1	1.0	1.2	1.5	1.7	1.6	1.5	1.2	1.1
1988	0.7	0.5	0.1	-0.3	-0.9	-1.3	-1.4	-1.2	-1.3	-1.6	-2.0	-2.0
1989	-1.8	-1.6	-1.2	-0.9	-0.7	-0.4	-0.4	-0.4	-0.4	-0.3	-0.2	-0.1
1990	0.1	0.1	0.3	0.3	0.2	0.2	0.3	0.3	0.3	0.3	0.3	0.4
1991	0.4	0.4	0.3	0.3	0.6	0.8	1.0	0.9	0.9	0.9	1.3	1.6
1992	1.8	1.7	1.5	1.4	1.2	0.9	0.5	0.2	-0.1	-0.1	0.1	0.3
1993	0.4	0.4	0.5	0.7	0.7	0.7	0.4	0.3	0.3	0.3	0.3	0.3
1994	0.2	0.2	0.2	0.3	0.4	0.4	0.5	0.5	0.7	0.9	1.3	1.3
1995	1.2	0.9	0.6	0.3	0.2	0.1	-0.1	-0.2	-0.5	-0.6	-0.8	-0.8
1996	-0.8	-0.7	-0.5	-0.3	-0.2	-0.2	-0.1	-0.2	-0.1	-0.2	-0.3	-0.4
1997	-0.4	-0.3	-0.1	0.3	0.8	1.3	1.7	2.0	2.2	2.4	2.5	2.5
1998	2.3	2.0	1.4	1.1	0.4	-0.1	-0.7	-1.0	-1.1	-1.2	-1.4	-1.5
1999	-1.5	-1.2	-0.9	-0.8	-0.8	-0.8	-0.9	-1.0	-1.0	-1.2	-1.4	-1.7
2000	-1.7	-1.4	-1.0	-0.8	-0.6	-0.6	-0.4	-0.4	-0.4	-0.5	-0.7	-0.7
2001	-0.7	-0.5	-0.4	-0.3	-0.1	0.1	0.1	0.0	0.0	-0.1	-0.1	-0.2



Pacific Niño 3.4 SST Outlook

A majority of ENSO forecasts indicate ENSO-neutral conditions will continue through Northern Hemisphere spring 2009.





Figure provided by the International Research Institute (IRI) for Climate and Society (updated 15 July 2008).




SST Outlook: NCEP CFS Forecast Issued 17 August 2008



The CFS ensemble mean (heavy blue line) predicts ENSO-neutral conditions through Northern Hemisphere Spring 2009.


Forecast *Nino3.4* SST anomalies from CFS






U. S. Seasonal Outlooks August – October 2008

Temperature



Precipitation



These seasonal outlooks combine long-term trends and soil moisture effects.



Summary

- **ENSO-neutral conditions are present in the equatorial Pacific Ocean.**
- **Equatorial SSTs in the central Pacific Ocean have returned to near-average, while positive SST anomalies continue in the eastern Pacific.**
- **Aspects of the atmospheric circulation and pattern of tropical convection reflect a lingering La Niña signal, particularly over the western and central Pacific.**
- **Based on recent SST trends and model forecasts, ENSO-neutral conditions are expected to continue through the Northern Hemisphere Fall 2008.**